

COMPLEXES OF MODULES OVER EXCEPTIONAL LIE SUPERALGEBRAS $E(3, 8)$ AND $E(5, 10)$.

VICTOR G. KAC* AND ALEXEI RUDAKOV

ABSTRACT. In this paper complexes of generalized Verma modules over the infinite-dimensional exceptional Lie superalgebras $E(3, 8)$ and $E(5, 10)$ are constructed and studied.

0. Introduction.

In our papers [KR1]–[KR3] we constructed all degenerate irreducible modules over the exceptional Lie superalgebra $E(3, 6)$. In the present paper we apply the same method to the exceptional Lie superalgebras $E(3, 8)$ and $E(5, 10)$.

The Lie superalgebra $E(3, 8)$ is strikingly similar to $E(3, 6)$. In particular, as in the case of $E(3, 6)$, the maximal compact subgroup of the group of automorphisms of $E(3, 8)$ is isomorphic to the group of symmetries of the Standard Model. However, as the computer calculations by Joris van Jeugt show, the fundamental particle contents in the $E(3, 8)$ case is completely different from that in the $E(3, 6)$ case [KR2]. All the nice features of the latter case, like the CPT symmetry, completely disappear in the former case. We believe that the main reason behind this is that, unlike $E(3, 6)$, $E(3, 8)$ cannot be embedded in $E(5, 10)$, which, we believe, is the algebra of symmetries of the SU_5 Grand Unified Model (the maximal compact subgroup of the automorphism group of $E(5, 10)$ is SU_5).

However, similarity with $E(3, 6)$ allows us to apply to $E(3, 8)$ all the arguments from [KR2] almost verbatim, and Figure 1 of the present paper that depicts all degenerate $E(3, 8)$ -modules is almost the same as Figure 3 from [KR2] for $E(3, 6)$.

The picture in the $E(5, 10)$ case is quite different (see Figure 2). We believe that it depicts all degenerate irreducible $E(5, 10)$ -modules, but we still do not have a proof.

1. Morphisms between generalized Verma modules.

Let $L = \bigoplus_{j \in \mathbb{Z}} \mathfrak{g}_j$ be a $\mathbb{Z}$ -graded Lie superalgebra by finite-dimensional vector spaces. Let

$$L_- = \bigoplus_{j < 0} \mathfrak{g}_j, \quad L_+ = \bigoplus_{j > 0} \mathfrak{g}_j, \quad L_0 = \mathfrak{g}_0 + L_+.$$

Given a $\mathfrak{g}_0$ -module V , we extend it to a L_0 -module by letting L_+ act trivially, and define the induced L -module

$$M(V) = U(L) \otimes_{U(L_0)} V.$$

If V is a finite-dimensional irreducible $\mathfrak{g}_0$ -module, the L -module $M(V)$ is called a *generalized Verma module* (associated to V), and it is called *degenerate* if it is not irreducible.

* Supported in part by NSF grant DMS-9970007.

Let A and B be two $\mathfrak{g}_0$ -modules and let $\text{Hom}(A, B)$ be the $\mathfrak{g}_0$ -module of linear maps from A to B . The following proposition will be extensively used to construct morphisms between the L -modules $M(A)$ and $M(B)$.

Proposition 1.1. *Let $\Phi \in M(\text{Hom}(A, B))$ be such that*

$$(1.1) \quad v \cdot \Phi = 0 \text{ for all } v \in L_0.$$

Then we can construct a well-defined morphism of L -modules

$$\varphi : M(A) \rightarrow M(B)$$

by the rule $\varphi(u \otimes a) = u \Phi(a)$. Explicitly, write $\Phi = \sum_m u_m \otimes \ell_m$, where $u_m \in U(L)$, $\ell_m \in \text{Hom}(A, B)$. Then

$$(1.2) \quad \varphi(u \otimes a) = \sum_m (u u_m) \otimes \ell_m(a).$$

Proof. We have to prove that for $v \in U(L_0)$,

$$\varphi(uv \otimes a) = \varphi(u \otimes va),$$

in order to conclude that φ is well-defined. Notice that condition (1.1) means

$$(1.3) \quad \sum_m [v, u_m] \otimes \ell_m + \sum_m u_m \otimes v \ell_m = 0.$$

Therefore we have:

$$\begin{aligned} \varphi(uv \otimes a) &= \sum_m uv u_m \otimes \ell_m(a) \\ &= \sum_m u[v, u_m] \otimes \ell_m(a) + \sum_m u u_m v \otimes \ell_m(a) \\ &= \sum_m u[v, u_m] \otimes \ell_m(a) + \sum_m u u_m \otimes v(\ell_m(a)) \\ &= \sum_m u[v, u_m] \otimes \ell_m(a) + \sum_m u u_m \otimes (v \ell_m)(a) + \sum_m u u_m \otimes \ell_m(va) \\ (\text{by (1.3) and (1.2)}) &= \sum_m u u_m \otimes \ell_m(va) = \varphi(u \otimes va). \end{aligned}$$

The fact that φ defines a morphism of L -modules is immediate from the definition. $\square$

Remark 1.1. If L_0 is generated by $\mathfrak{g}_0$ and a subset of $T \subset L_+$, then condition (1.1) is equivalent to

$$(1.4a) \quad \mathfrak{g}_0 \cdot \Phi = 0$$

$$(1.4b) \quad a \cdot \Phi = 0 \text{ for all } a \in T.$$

Condition (1.4a) usually gives a hint to a possible shape of Φ and is checked by general invariant-theoretical considerations. After that (1.4b) is usually checked by a direct calculation.

Remark 1.2. We can view $M(V)$ also as the induced $(L_- \oplus \mathfrak{g}_0)$ -module: $U(L_- \oplus \mathfrak{g}_0) \otimes_{U(\mathfrak{g}_0)} V$. Then condition (1.4a) on $\Phi = \sum_m u_m \otimes \ell_m$, where $u_m \in U(L_- \oplus \mathfrak{g}_0)$, $\ell_m \in \text{Hom}(A, B)$, suffices in order (1.2) to give a well-defined morphism of $(L_- \oplus \mathfrak{g}_0)$ -modules. One can also replace $\mathfrak{g}_0$ by any of its subalgebras.

2. Lie superalgebras $E(3, 6)$, $E(3, 8)$ and $E(5, 10)$.

Recall some standard notation:

$$W_n = \left\{ \sum_{j=1}^n P_j(x) \partial_j \mid P_j \in \mathbb{C}[[x_1, \dots, x_n]], \partial_j \equiv \partial / \partial x_j \right\}$$

denotes the Lie algebra of formal vector fields in n indeterminates:

$$S_n = \left\{ D = \sum_i P_i \partial_i \mid \operatorname{div} D \equiv \sum_i \partial_i P_i = 0 \right\}$$

denotes the Lie subalgebra of divergenceless formal vector fields; $\Omega^k(n)$ denotes the associative algebra of formal differential forms of degree k in n indeterminates, $\Omega_{cl}^k(n)$ denoted the subspace of closed forms.

The Lie algebra W_n acts on $\Omega^k(n)$ via Lie derivative $D \rightarrow L_D$. Given $\lambda \in \mathbb{C}$ one can twist this action:

$$D\omega = L_D \omega + \lambda(\operatorname{div} D)\omega.$$

The W_n -module thus obtained is denoted by $\Omega^k(n)^\lambda$. Recall the following obvious isomorphism of W_n -modules

$$(2.1) \quad \Omega^0(n) \simeq \Omega^n(n)^{-1},$$

and the following slightly less obvious isomorphism of W_n -modules

$$(2.2) \quad W_n \simeq \Omega^{n-1}(n)^{-1}.$$

The latter is obtained by mapping a vector field $D \in W_n$ to the $(n-1)$ -form $\iota_D(dx_1 \wedge \dots \wedge dx_n)$. Note that (2.2) induces an isomorphism of S_n -modules:

$$(2.3) \quad S_n \simeq \Omega_{cl}^{n-1}(n).$$

Recall that the Lie superalgebra $E(5, 10) = E(5, 10)_0 + E(5, 10)_1$ is constructed as follows [K], [CK]:

$$E(5, 10)_0 = S_5, \quad E(5, 10)_1 = \Omega_{cl}^2(5),$$

$E(5, 10)_0$ acts on $E(5, 10)_1$ via the Lie derivative, and $[\omega_2, \omega'_2] = \omega_2 \wedge \omega'_2 \in \Omega_{cl}^4(5) = S_5$ (see (2.3)) for $\omega_2, \omega'_2 \in E(5, 10)_1$.

Next, recall the construction of the Lie superalgebras $E^\flat := E(3, 6)$ and $E^\sharp := E(3, 8)$ [CK]:

$$\begin{aligned} E_0^\flat &= E_0^\sharp = W_3 + s\ell_2(\Omega^0(3)) \quad (\text{the natural semidirect sum}); \\ E_1^\flat &= \Omega^1(3)^{-1/2} \otimes \mathbb{C}^2, \quad E_1^\sharp = (\Omega^0(3)^{-1/2} \otimes \mathbb{C}^2) + (\Omega^2(3)^{-1/2} \otimes \mathbb{C}^2). \end{aligned}$$

The action of the even on the odd parts is defined via the Lie derivative and the multiplication of a function and a differential form. The bracket of two odd elements is defined by using the identifications (2.1) and (2.2) as follows. For $\omega_i, \omega'_i \in \Omega^i(3)$ and $v, v' \in \mathbb{C}^2$ one defines the following bracket of two elements from $E_1^\flat$:

$$(2.4) \quad [\omega_1 \otimes v, \omega'_1 \otimes v'] = -(\omega_1 \wedge \omega'_1) \otimes (v \wedge v') - (d\omega_1 \wedge \omega'_1 + \omega_1 \wedge d\omega'_1) \otimes (v \cdot v'),$$

and the following bracket of two elements from $E_1^\sharp$:

$$(2.5) \quad \begin{aligned} [\omega_2 \otimes v, \omega'_2 \otimes v'] &= 0, \quad [\omega_0 \otimes v, \omega'_0 \otimes v'] = -(d\omega_0 \wedge d\omega'_0) \otimes (v \wedge v'), \\ [\omega_0 \otimes v, \omega_2 \otimes v'] &= -(\omega_0 \wedge \omega_2) \otimes (v \wedge v') - (d\omega_0 \wedge \omega_2 - \omega_0 \wedge d\omega_2) \otimes (v \cdot v'). \end{aligned}$$

Recall also an embedding of E^b in $E(5, 10)$ [CK], [KR2]. For that let $z_+ = x_4$, $z_- = x_5$, $\partial_+ = \partial_4$, $\partial_- = \partial_5$, and let ϵ^+, ϵ^- denote the standard basis of $\mathbb{C}^2$. Then E_0^b is embedded in $E(5, 10)_0 = S_5$ by $(D \in W_3, a, b, c \in \Omega^0(3))$:

(2.6)

$$D \mapsto D - \frac{1}{2}(\operatorname{div} D)(z_+ \partial_+ + z_- \partial_-), \quad \begin{pmatrix} a & b \\ c & -a \end{pmatrix} \mapsto a(z_+ \partial_+ - z_- \partial_-) + bz_+ \partial_- + cz_- \partial_+,$$

and E_1^b is embedded in $E(5, 10)_1 = \Omega_{\text{cl}}^2(5)$ by $(f \in \Omega^0(3))$:

(2.7)

$$f dx_i \otimes \epsilon^\pm \mapsto z_\pm dx_i \wedge df + f dx_i \wedge dz_\pm.$$

Introduce the following subalgebras $S^b \subset E^b$ and $S^\sharp \subset E^\sharp$:

$$\begin{aligned} S_0^b &= S_0^\sharp = W_3 + \mathbb{C} \otimes \mathfrak{sl}_2(\mathbb{C}), \\ S_1^b &= \Omega_{\text{cl}}^1(3)^{-1/2} \otimes \mathbb{C}^2, \quad S_1^\sharp = \Omega^0(3)^{-1/2} \otimes \mathbb{C}^2. \end{aligned}$$

Proposition 2.1. *The map $S^\sharp \rightarrow S^b$, which is identical on $S_0^\sharp$ and sends $f \otimes v \in S_1^\sharp$ to $df \otimes v \in S_1^b$ is a surjective homomorphism of Lie superalgebras with 2-dimensional central kernel $\mathbb{C} \otimes \mathbb{C}^2 \subset S_1^\sharp$.*

Proof. It is straightforward using (2.4) and (2.5). □

This proposition is probably the main reason for a remarkable similarity between representation theories of $E^\sharp$ and E^b . We shall stress this similarity in our notation and develop representation theory of $E^\sharp$ along the same lines as that of E^b done in [KR1], [KR2]. Sometimes we shall drop the superscript b or $\sharp$ when the situation is the same.

Recall that E^b carries a unique irreducible consistent $\mathbb{Z}$ -gradation. It has depth 2, and it is defined by:

$$(2.8) \quad \deg x_i = -\deg \partial_i = 2, \quad \deg \partial_i = -1, \quad \deg \epsilon^\pm = 0, \quad \deg \mathfrak{sl}_2(\mathbb{C}) = 0.$$

The “non-positive” part of this $\mathbb{Z}$ -gradation is as follows:

$$\begin{aligned} \mathfrak{g}_{-2} &= \langle \partial_i \mid i = 1, 2, 3 \rangle, \quad \mathfrak{g}_{-1} = \langle d_i^a := \epsilon^a \otimes dx_i = dx_i \wedge dz_a \mid i = 1, 2, 3, a = +, - \rangle, \\ \mathfrak{g}_0 &= \mathfrak{sl}_3(\mathbb{C}) \oplus \mathfrak{sl}_2(\mathbb{C}) \oplus CY, \end{aligned}$$

where

$$(2.9) \quad \mathfrak{sl}_3(\mathbb{C}) = \langle h_1 = x_1 \partial_1 - x_2 \partial_2, h_2 = x_2 \partial_2 - x_3 \partial_3, e_1 = x_1 \partial_2, \\ e_2 = x_2 \partial_3, e_{12} = x_1 \partial_3, f_1 = x_2 \partial_1, f_2 = x_3 \partial_2, f_{12} = x_3 \partial_1 \rangle,$$

$$(2.10) \quad \mathfrak{sl}_2(\mathbb{C}) = \langle h_3 = z_+ \partial_+ - z_- \partial_-, e_3 = z_+ \partial_-, f_3 = z_- \partial_+ \rangle,$$

$$(2.11) \quad Y = \frac{2}{3} \sum x_i \partial_i - (z_+ \partial_+ + z_- \partial_-).$$

The eigenspace decomposition of $\operatorname{ad}(3Y)$ coincides with the consistent $\mathbb{Z}$ -grading of E^b . We fix the Cartan subalgebra $\mathcal{H} = \langle h_1, h_2, h_3, Y \rangle$ and the Borel subalgebra $\mathcal{B} = \mathcal{H} + \langle e_i, (i = 1, 2, 3), e_{12} \rangle$ of $\mathfrak{g}_0$. Then $f_0 := d_1^+$ is the highest weight vector of the (irreducible) $\mathfrak{g}_0$ -module $\mathfrak{g}_{-1}$, the vectors

$$e'_0 := x_3 d_3^- \quad \text{and} \quad e_0^b := x_3 d_2^- - x_2 d_3^- + 2z_- dx_2 \wedge dx_3$$

are all the lowest weight vectors of the $\mathfrak{g}_0$ -module $\mathfrak{g}_1$, and one has:

$$(2.12) \quad [e'_0, f_0] = f_2,$$

$$(2.13) \quad [e_0^\flat, f_0] = \frac{2}{3}h_1 + \frac{1}{3}h_2 - h_3 - Y =: h_0^\flat,$$

so that

$$h_0^\flat = -x_2\partial_2 - x_3\partial_3 + 2z_-\partial_-.$$

The following relations are also important to keep in mind:

$$(2.14) \quad [e'_0, d_1^+] = f_2, \quad [e'_0, d_2^+] = -f_{12}, \quad [e'_0, d_3^+] = 0, \quad [e'_0, d_i^-] = 0,$$

$$(2.15) \quad [d_i^\pm, d_j^\pm] = 0, \quad [d_i^+, d_j^-] + [d_j^+, d_i^-] = 0.$$

Recall that $\mathfrak{g}_0$ along with the elements $f_0, e_0^\flat, e'_0$ generate the Lie superalgebra $E^\flat$ [CK].

The Lie superalgebra $E^\sharp$ carries a unique consistent irreducible $\mathbb{Z}$ -gradation of depth 3:

$$E^\sharp = \bigoplus_{j \geq -3} \mathfrak{g}_j.$$

It is defined by:

$$(2.16) \quad \deg x_i = -\deg \partial_i = \deg dx_i = 2, \quad \deg \epsilon^\pm = -3, \quad \deg s\ell_2(\mathbb{C}) = 0.$$

In view of Proposition 2.1 and the above $E^\flat$ -notation, we introduce the following $E^\sharp$ -notation:

$$(2.17) \quad \begin{aligned} d^\pm &:= 1 \otimes \epsilon^\pm, \quad d_i^\pm := x_i \otimes \epsilon^\pm, \quad e'_0 := \frac{1}{2}x_3^2 \otimes \epsilon^\pm, \quad f_0 := d_1^+, \\ e_0^\sharp &:= -(dx_2 \wedge dx_3) \otimes \epsilon^-, \quad h_0^\sharp := \frac{2}{3}h_1 + \frac{1}{3}h_2 - \frac{1}{2}h_3 + \frac{1}{2}Y. \end{aligned}$$

If, in analogy with $E^\flat$, we denote $\epsilon^a = dz_a$, then $\mathfrak{g}_{-3} = \langle dz_+, dz_- \rangle$, $\mathfrak{g}_{-2}$, $\mathfrak{g}_{-1}$ and $\mathfrak{g}_0$ are the same as for $E^\flat$ (except that now $[\mathfrak{g}_{-1}, \mathfrak{g}_{-2}] \neq 0$), and the relations (2.12), (2.13), (2.14), (2.15) still hold, but the formula for h_0 is different:

$$(2.18) \quad h_0^\sharp = \frac{2}{3}h_1 + \frac{1}{3}h_2 - \frac{1}{2}h_3 + \frac{1}{2}Y = x_1\partial_1 - z_+\partial_+.$$

As in the $E^\flat$ case, the elements e_i, f_i, h_i for $i = 0, 1, 2, 3$ along with e'_0 generate $E^\sharp$, the elements $e_0^\sharp$ and e'_0 are all lowest weight vectors of the $\mathfrak{g}_0$ -module $\mathfrak{g}_1$, and $\mathfrak{g}_0$ along with e'_0 and $e_0^\sharp$ generate the subalgebra $\bigoplus_{j \geq 0} \mathfrak{g}_j$ [CK]. Thus, by Remark 1.1, condition (1.4b) is equivalent to

$$(2.19) \quad e'_0 \cdot \Phi = 0 \quad \text{and} \quad e_0^\sharp \cdot \Phi = 0.$$

3. Complexes of degenerate Verma modules over $E(3, 6)$.

Let W be a finite-dimensional symplectic vector space and let H be the corresponding Heisenberg algebra

$$H = T(W)/(v \cdot w - w \cdot v - (v, w) \cdot 1)$$

where $T(W)$ denotes the tensor algebra over W and $(\cdot, \cdot)$ is the non-degenerate symplectic form on W . Given two transversal Lagrangian subspaces $L, L' \subset W$, we have a canonical isomorphism of symplectic spaces: $W = L + L' \simeq L \oplus L^*$, and we can canonically identify the symmetric algebra $S(L)$ with the factor of H by the left ideal generated by L' :

$$(3.1) \quad V_L := H/(L') \simeq S(L)\mathbf{1}_L, \quad \text{where } \mathbf{1}_L = 1 + L'.$$

One thus acquires an H -module structure on $S(L)$.

We construct a symplectic space W by taking $x_1, x_2, x_3, z_+, z_-, \partial_1, \partial_2, \partial_3, \partial_+, \partial_-$ as a basis with the first half being dual to the second half:

$$(3.2) \quad (\partial_i, x_j) = -(x_j, \partial_i) = \delta_{ij}, \quad (\partial_a, z_b) = -(z_b, \partial_a) = \delta_{a,b}, \quad \text{all other pairings zero.}$$

In general the decomposition $W = L + L' \simeq L \oplus L^*$ provides the canonical maps:

$$\text{End}(L) \xrightarrow{\sim} L \otimes L^* \xrightarrow{\sim} L \cdot L' \hookrightarrow H,$$

which induce a Lie algebra homomorphism: $g\ell(L) \rightarrow H_{Lie}$, where the Lie algebra structure is defined by the usual commutator.

We consider the following subspaces of W :

$$(3.3) \quad \begin{aligned} L_A &= L'_D = \langle x_i, z_a \rangle, & L_B &= L'_C = \langle x_i, \partial_a \rangle, \\ L_C &= L'_B = \langle \partial_c, z_a \rangle, & L_D &= L'_A = \langle \partial_i, \partial_a \rangle, \end{aligned}$$

where $i = 1, 2, 3$, $a = +, -$. Note that these are the only $\mathfrak{g}_0$ -invariant Lagrangian subspaces of W .

As formulae (2.9)–(2.11) determine the inclusion $\mathfrak{g}_0 \hookrightarrow g\ell(V_X)$, where $X = A, B, C$ or D , we get a Lie algebra monomorphism:

$$(3.4) \quad \mathfrak{g}_0 \hookrightarrow g\ell(V_X) \hookrightarrow H_{Lie}.$$

Thus we get a $\mathfrak{g}_0$ -action on V_X . Let us notice that by (3.4)

$$(3.5) \quad Y \rightarrow Y^\flat = \frac{2}{3} \left(\sum_i x_i \partial_i \right) - \left(\sum_a z_a \partial_a \right)$$

and

$$Y^\flat \mathbf{1}_A = 0, \quad Y^\flat \mathbf{1}_B = 2 \mathbf{1}_B, \quad Y^\flat \mathbf{1}_C = -2 \mathbf{1}_C, \quad Y^\flat \mathbf{1}_D = 0,$$

as it should be for $E^\flat = E(3, 6)$ (see [KR2]).

In the $E(3, 8)$ case we modify the $\mathfrak{g}_0$ -action on V_X leaving it the same for $sl_3(\mathbb{C}) \oplus sl_2(\mathbb{C}) \subset \mathfrak{g}_0$, but letting

$$(3.6) \quad Y \mapsto Y^\sharp = -\frac{4}{3} \left(\sum_i x_i \partial_i \right) + \left(\sum_a z_a \partial_a \right) = Y^\flat + 2T, \quad \text{where } T = -\sum_i x_i \partial_i + \sum_a z_a \partial_a.$$

We have got:

$$\begin{aligned} Y x_i^p z_a^r \mathbf{1}_A &= \left(-\frac{4}{3}p + r \right) x_i^p z_a^r \mathbf{1}_A, \\ Y x_i^p \partial_a^r \mathbf{1}_B &= \left(\frac{4}{3}p - r - 2 \right) x_i^p \partial_a^r \mathbf{1}_B, \\ Y \partial_i^q z_a^r \mathbf{1}_C &= \left(\frac{4}{3}q + r + 4 \right) \partial_i^q z_a^r \mathbf{1}_C, \\ Y \partial_i^q \partial_a^r \mathbf{1}_D &= \left(\frac{4}{3}q - r + 2 \right) \partial_i^q \partial_a^r \mathbf{1}_D. \end{aligned}$$

Let $F(p, q; r; y)$ denote the finite-dimensional irreducible $\mathfrak{g}_0$ -module with highest weight $(p, q; r; y)$, where $p, q, r \in \mathbb{Z}_+$, $y \in \mathbb{C}$, and let $M(p, q; r; y) = M(F(p, q; r; y))$ be the corresponding generalized Verma module over $E(3, 8)$. This module has a unique irreducible quotient denoted by $I(p, q; r; y)$. The latter module is called degenerate if the former is.

We announce below a classification of all degenerate irreducible $E(3, 8)$ -modules.

Theorem 3.1. *All irreducible degenerate $E(3, 8)$ -modules $I(p, q; r, y)$ are as follows $(p, q, r \in \mathbb{Z}_+)$:*

$$\begin{aligned} \text{type } A: & \quad I(p, 0; r; y_A), \quad y_A = -\frac{4}{3}p + r; \\ \text{type } B: & \quad I(p, 0; r; y_B), \quad y_B = -\frac{4}{3}p - r - 2; \\ \text{type } C: & \quad I(0, q; r; y_C), \quad y_C = \frac{4}{3}q + r + 4; \\ \text{type } D: & \quad I(0, q; r; y_D), \quad y_D = \frac{4}{3}q - r + 2, \text{ and } (q, r) \neq (0, 0), . \end{aligned}$$

We shall construct below certain $E(3, 8)$ -morphisms between the modules $M(p, q; r, y_X)$. This will imply that all modules $I(p, q; r, y)$ on the list are degenerate. The proof of the fact that the list is complete will be published elsewhere.

The theorem means that all the degenerate generalized Verma modules over $E(3, 8)$ are in fact the direct summands of induced modules $M(V_X)$, $X = A, B, C, D$:

$$\begin{aligned} M(V_X) &= \bigoplus_{m, n \in \mathbb{Z}} M(V_X^{m, n}), \text{ where} \\ V_X^{m, n} &= \left\{ f \mathbf{1}_X \mid \left(\sum x_i \partial_i \right) f = m f, \left(\sum z_a \partial_a \right) f = n f \right\}, \end{aligned}$$

(we normalize degree of $\mathbf{1}_X$ as $(0, 0)$).

We construct morphisms between these modules with the help of Proposition 1.1. As in the E^b case [KR2], introduce the following operators on $M(\text{Hom}(V_X, V_X))$:

$$\begin{aligned} \nabla &= \Delta^+ \delta_+ + \Delta^- \delta_- = \delta_1 \partial_1 + \delta_2 \partial_2 + \delta_3 \partial_3, \text{ where} \\ \Delta^\pm &= \sum_{i=1}^3 d_i^\pm \otimes \partial_i, \quad \delta_i = \sum_{\alpha=\pm} d_i^\alpha \otimes \partial_\alpha. \end{aligned}$$

Proposition 3.2. (a) *The element ∇ gives a well-defined morphism $M(V_X) \rightarrow M(V_X)$, $X = A, B, C, D$, by formula (1.2).*

(b) $\nabla^2 = 0$.

Proof. The proof of (b) is the same as in [KR2]. In order to prove (a), we have to check conditions (1.4a) and (1.4b). It is obvious that $\Delta^\pm$ (resp. δ_i) are $sl_3(\mathbb{C})$ -(resp. $sl_2(\mathbb{C})$)-invariant. Using both formulas for ∇ , we conclude that it is $\mathfrak{g}_0$ -invariant, proving (1.4a). In order to check (1.4b), first note that

$$e'_0 \nabla = (f_2 \partial_1 \partial_+ - f_{12} \partial_2 \partial_+) = (x_3 \partial_2 \partial_1 - x_3 \partial_1 \partial_2) \partial_+ = 0.$$

Now

$$\begin{aligned} e_0^\# \nabla &= h_0^\# \partial_1 \partial_+ + f_1 \partial_2 \partial_+ + f_{12} \partial_3 \partial_1 - f_3 \partial_1 \partial_- \\ &= (x_1 \partial_1 - z_+ \partial_+ + T) \partial_1 \partial_+ + x_2 \partial_1 \partial_2 \partial_+ + x_3 \partial_1 \partial_3 \partial_+ - z_- \partial_+ \partial_1 \partial_- \\ &= (x_1 \partial_1 - z_+ \partial_+ + T + x_2 \partial_2 + x_3 \partial_3 - z_- \partial_-) \partial_1 \partial_+ = 0, \end{aligned}$$

where we use (3.6) to check that

$$(3.7) \quad h_0^\# = \frac{2}{3}h_1 + \frac{1}{3}h_2 - \frac{1}{2}h_3 + \frac{1}{2}Y^\# = x_1 \partial_1 - z_+ \partial_+ + T.$$

□

Let $M_{X'} = M(V_{X'})$, where:

$$\begin{aligned} (3.8) \quad V_{A'} &= \mathbb{C}[x_1, x_2, x_3] \mathbf{1}_A = \oplus_{p \geq 0} V_A^{p, 0}, \quad V_{B'} = \mathbb{C}[x_1, x_2, x_3] \mathbf{1}_B = \oplus_{p \geq 0} V_B^{p, 0} \\ V_{C'} &= \mathbb{C}[\partial_1, \partial_2, \partial_3] \mathbf{1}_C = \oplus_{q \geq 0} V_C^{-q, 0}, \quad V_{D'} = \mathbb{C}[\partial_1, \partial_2, \partial_3] \mathbf{1}_D = \oplus_{q \geq 0} V_D^{-q, 0} \end{aligned}$$

and let

$$(3.9) \quad \nabla_2 = \Delta^- \Delta^+ = d_1^- \Delta^+ \partial_1 + d_2^- \Delta^+ \partial_2 + d_3^- \Delta^+ \partial_3.$$

We define the morphism $\nabla_2 : M_{A'} \rightarrow M_{B'}$ by extending the map defined by ∇_2 as follows

$$(3.10) \quad \begin{array}{ccc} V_{A'} \xrightarrow{\sim} \mathbb{C}[x_1, x_2, x_3] & \longrightarrow & U(L_-) \otimes \mathbb{C}[x_1, x_2, x_3] \mathbf{1}_B \simeq U(L_-) \otimes V_{B'}, \\ f \mathbf{1}_A & \mapsto & f \quad \mapsto \quad \nabla_2 f \mathbf{1}_B. \end{array}$$

In order to apply Proposition 1.1, we have to check conditions (1.4a) and (1.4b) for ∇_2 . As before, condition (1.4a) obviously holds. Now

$$e'_0 \nabla_2 \mathbf{1}_B = - \sum_i d_i^- (f_2 \partial_1 \partial_- - f_{12} \partial_2) \partial_i f \mathbf{1}_B = 0$$

because $f_2 \partial_1 - f_{12} \partial_2 = x_3 \partial_2 \partial_1 - x_3 \partial_1 \partial_2 = 0$. Furthermore:

$$\begin{aligned} e_0^\# \nabla_2 f \mathbf{1}_B &= \left(- \sum_i d_i^- (h_0^\# \partial_1 + f_1 \partial_2 + f_{12} \partial_3) \partial_i - f_3 \Delta^+ \partial_1 \right) f \mathbf{1}_B \\ &= - \left(\sum_i d_i^- (x_1 \partial_1 - z_+ \partial_+ + T + x_2 \partial_2 + x_3 \partial_3) \partial_i \partial_- - \Delta^- \partial_1 - \Delta^+ \partial_1 f_3 \right) f \mathbf{1}_B. \end{aligned}$$

As $z_\pm \partial_\pm f \mathbf{1}_B = f z_\pm \partial_\pm \mathbf{1}_B = f(-1 + \partial_\pm z_\pm) \mathbf{1}_B = -f \mathbf{1}_B$, and $f_3 f \mathbf{1}_B = 0$, we conclude that

$$\begin{aligned} -e_0^\# \nabla_2 f \mathbf{1}_B &= \sum_i d_i^- (x_1 \partial_1 + x_2 \partial_2 + x_3 \partial_3 - z_+ \partial_+ + T + 1) f \mathbf{1}_B \\ &= \sum_i d_i^- (z_- \partial_- + 1) f \mathbf{1}_B = 0. \end{aligned}$$

Thus Proposition 1.1 applies and we get the morphism $\nabla_2 : M_{A'} \rightarrow M_{B'}$.

In exactly the same fashion we construct the morphism $\nabla_2 : M_{C'} \rightarrow M_{D'}$ by taking $f \in \mathbb{C}[\partial_1, \partial_2, \partial_3]$.

Thus we have proved part (a) of the following proposition; part (b) was checked in [KR2].

Proposition 3.3. (a) *Formulae (3.9) and (3.10) define the morphisms $\nabla_2 : M_{A'} \rightarrow M_{B'}$, $\nabla_2 : M_{C'} \rightarrow M_{D'}$ of $E(3, 8)$ -modules.*

(b) $\nabla \nabla_2 = 0$, $\nabla_2 \nabla = 0$.

Similarly we can construct morphisms with the help of the element

$$(3.11) \quad \nabla_3 = \delta_1 \delta_2 \delta_3 = \sum_{a,b,c} d_1^a d_2^b d_3^c \otimes \partial_a \partial_b \partial_c, \quad (a, b, c = +, -).$$

Let us consider modules $M_{X''} = M(V_{X''})$, where

$$\begin{aligned} V_{A''} &= \mathbb{C}[z_+, z_-] \mathbf{1}_A = \oplus_{r \geq 0} V_A^{0,r}, \quad V_{B''} = \mathbb{C}[\partial_+, \partial_-] \mathbf{1}_B = \oplus_{r \geq 0} V_B^{0,-r}, \\ V_{C''} &= \mathbb{C}[z_+, z_-] \mathbf{1}_C = \oplus_{r \geq 0} V_C^{0,r}, \quad V_{D''} = \mathbb{C}[\partial_+, \partial_-] \mathbf{1}_D = \oplus_{r \geq 0} V_D^{0,-r}. \end{aligned}$$

We construct morphisms $\nabla_3 : M_{A''} \rightarrow M_{C''}$ and $\nabla_3 : M_{B''} \rightarrow M_{D''}$ by extending the maps $\nabla_3' : V_{A''} \rightarrow U(L_-) \otimes V_{C''}$, $\nabla_3'' : V_{B''} \rightarrow U(L_-) \otimes V_{D''}$ given by the left and right diagrams below:

$$(3.12) \quad \begin{array}{ccc} \mathbb{C}[z_a] \cdot \mathbf{1}_A \xrightarrow{\sim} & \mathbb{C}[z_+, z_-] & \mathbb{C}[\partial_a] \cdot \mathbf{1}_B \xrightarrow{\sim} & \mathbb{C}[\partial_+, \partial_-] \\ & \downarrow \nabla_3 & & \downarrow \nabla_3 \\ & U(L_-) \otimes (\mathbb{C}[z_+, z_-] \mathbf{1}_C) & & U(L_-) \otimes (\mathbb{C}[\partial_+, \partial_-] \mathbf{1}_D). \end{array}$$

Here the horizontal maps are naturally $sl_3(\mathbb{C}) \oplus sl_2(\mathbb{C})$ -isomorphisms, but we have to define the action of Y on the target demanding the map to be a $\mathfrak{g}_0$ -isomorphism. With this in mind we have to check that Y commutes with ∇_3 .

We shall use the Einstein summation convention argument for the vertical maps ∇_3 given by (3.12). Then for $f \in \mathbb{C}[z_+, z_-]$ we have $Y(f\mathbf{1}_A) = (\deg f)f\mathbf{1}_A$ and

$$\begin{aligned} Y(d_1^a d_2^b d_3^c \otimes \partial_a \partial_b \partial_c f \mathbf{1}_C) &= d_1^a d_2^b d_3^c \otimes (-1 + Y^\sharp) \partial_1 \partial_b \partial_c f \mathbf{1}_C \\ &= d_1^a d_2^b d_3^c \otimes \partial_a \partial_b \partial_c f (-4 + \deg f + Y^\sharp) \mathbf{1}_C \\ &= (\deg f) \cdot d_1^a d_2^b d_3^c \otimes \partial_a \partial_b \partial_c f \mathbf{1}_C. \end{aligned}$$

Similarly for $f \in \mathbb{C}[\partial_+, \partial_-]$, we have $Y(f\mathbf{1}_B) = (-\deg f - 2)f\mathbf{1}_B$ and

$$\begin{aligned} Y(d_1^a d_2^b d_3^c \otimes \partial_a \partial_b \partial_c f \mathbf{1}_D) &= d_1^a d_2^b d_3^c \otimes \partial_a \partial_b \partial_c f (-4 - \deg f + Y^\sharp) \mathbf{1}_D \\ &= (-4 - \deg f + 2) d_1^a d_2^b d_3^c \otimes \partial_a \partial_b \partial_c f \mathbf{1}_D, \end{aligned}$$

where $Y^\sharp$ defined by (3.6). Thus we get the commutativity.

One meets no problem checking $e'_0 \nabla_3 = 0$, but we consider calculations for $e_0^\sharp \nabla_3$ in more detail. If $f \in \mathbb{C}[z_+, z_-]$, then

$$e_0(\nabla_3 f \mathbf{1}_C) = h_0^\sharp d_2^b d_3^c \otimes \partial_+ \partial_b \partial_c f \mathbf{1}_C - f_3 d_2^b d_3^c \otimes \partial_- \partial_b \partial_c f \mathbf{1}_C,$$

because $d_1^a(x_2 \partial_1) d_3^c \otimes \partial_a \partial_+ \partial_c f \mathbf{1}_C = d_1^a d_2^b(x_3 \partial_1) \otimes \partial_a \partial_b \partial_+ f \mathbf{1}_C = 0$. Now $f_3(d_2^b d_3^c \otimes \partial_b \partial_c) = (d_2^b d_3^c \otimes \partial_b \partial_c) f_3$ and $h_0^\sharp(d_2^b d_3^c \otimes \partial_b \partial_c) = (d_2^b d_3^c \otimes \partial_b \partial_c)(h_0^\sharp - 2)$, where $h_0^\sharp = \frac{2}{3}h_1 + \frac{1}{3}h_2 - \frac{1}{2}h_3 + \frac{1}{2}Y^\sharp$, and again $Y^\sharp$ is defined by (3.6). Therefore

$$\begin{aligned} e_0^\sharp(\nabla_3 f \mathbf{1}_C) &= (d_2^b d_3^c \otimes \partial_b \partial_c)(h_0^\sharp \partial_+ - 2\partial_+ - z_+ \partial_+ \partial_-) f \mathbf{1}_C \\ &= (d_2^b d_3^c \otimes \partial_b \partial_c)(h_0^\sharp - z_- \partial_- - 2) \partial_+ f \mathbf{1}_C \\ &= (d_2^b d_3^c \otimes \partial_b \partial_c)(-x_2 \partial_2 - x_3 \partial_3 - 2) \partial_+ f \mathbf{1}_C \\ &= (d_2^b d_3^c \otimes \partial_b \partial_c)(-\partial_2 x_2 - \partial_3 x_3) \partial_+ f \mathbf{1}_C = 0. \end{aligned}$$

The calculations in the case $f \in \mathbb{C}[\partial_+, \partial_-]$ are very much the same. So we have proved part (a) of the following proposition; part (b) was checked in [KR2]

Proposition 3.4. (a) *Formulae (3.11) and (3.12) define the morphisms $\nabla_3 : M_{A''} \rightarrow M_{D''}$ and $M_{B''} \rightarrow M_{D''}$ of $E(3, 8)$ -modules.*

(b) $\nabla \cdot \nabla_3 = 0$, $\nabla_3 \cdot \nabla = 0$.

Furthermore, there are $E(3, 8)$ -module morphisms

$$\begin{aligned} \nabla'_4 &: M(00; 2; y_A) \rightarrow M(01; 1; y_D) \text{ and} \\ \nabla''_4 &: M(10; 0; y_A) \rightarrow M(00; 2; y_D) \end{aligned}$$

defined by formulae (2.14) and (2.17) from [KR2], applied to $E(3, 8)$. Arguments similar to those in [KR2] show that these are indeed well-defined morphisms.

Thus far we have constructed $E(3, 8)$ -homomorphisms ∇ , ∇_2 , ∇_3 , ∇'_4 , and ∇''_4 between generalized Verma modules. Note that these maps have degree 1, 2, 3 and 4, respectively with respect to the $\mathbb{Z}$ -gradation of $U(L_-)$ induced by that of $E(3, 8)$.

As in the case of $E(3, 6)$ [KR2], all these maps are illustrated in Figure 1. The nodes in the quadrants A, B, C, D represent generalized Verma modules $M(p, 0; r; y_X)$ if $X = A$ or B , and $M(0, q; r; y_X)$, if $X = C$ or D . The plain arrows represent ∇ , the dotted arrows represent ∇_2 , the interrupted arrows represent ∇_3 and the bold arrows represent ∇'_4 and ∇''_4 .




FIGURE 1.

Note that the generalized Verma modules $M(00; 1; y_A)$ and $M(00; 1; y_D)$ are isomorphic since $y_A = y_D = 1$. We shall identify them. This allows us to construct the $E(3, 8)$ -module homomorphism

$$\tilde{\nabla} : M(00; 1; y_A) \rightarrow M(01; 2; y_D),$$

which is NOT represented in Figure 1.

Note that $I(00; 1; y_A) = I(00; 1; y_D)$ is the coadjoint $E(3, 8)$ -module. It follows from the above propositions that if we remove the module $M(00; 1; y_D)$ from Figure 1, and draw $\tilde{\nabla}$ then all sequences in the modified Figure 1 become complexes. We denote by $H_A^{p, -r}$, $H_B^{p, r}$, $H_C^{-q, r}$ and $H_D^{-q, -r}$ the homology of these complexes at the position of $M(pq; r; y_X)$, $X = A, B, C, D$.

Theorem 3.5. (a) The kernels of all maps ∇ , ∇_2 , ∇_3 , ∇'_4 , ∇''_4 , $\tilde{\nabla}$ are maximal submodules.

(b) The homology $H_X^{m,n}$ is zero except for six cases listed (as $E(3, 8)$ -modules) below:

$$\begin{aligned} H_A^{0,0} &= \mathbb{C}, & H_A^{1,1} &= I(10; 0; -\frac{4}{3}), \\ H_A^{1,0} &= H_D^{0,-2} = I(00; 0; -2), \\ H_D^{-1,-1} &= H_D^{-1,-2} = I(00; 1; 1) \oplus \mathbb{C}. \end{aligned}$$

The proof is similar to that of the analogous $E(3, 6)$ -result in [KR2]. Note that this theorem gives the following explicit construction of all degenerate irreducible $E(3, 8)$ -modules:

$$I(pq; r; y_X) = M(pq; r; y_X) / \text{Ker } \nabla,$$

where ∇ is the corresponding map in the modified Figure 1.

4. Three series of degenerate Verma modules over $E(5, 10)$

As in [KR2] and in § 2, we use for the odd elements of $E(5, 10)$ the notation $d_{ij} = dx_i \wedge dx_j$ ($i, j = 1, 2, \dots, 5$); recall that we have the following commutation relation ($f, g \in \mathbb{C}[[x_1, \dots, x_5]]$):

$$[fd_{jk}, gd_{\ell m}] = \epsilon_{ijklm} \partial_i,$$

where ϵ_{ijklm} is the sign of the permutation $ijklm$ if all indices are distinct and 0 otherwise.

Recall that the Lie superalgebra $E(5, 10)$ carries a unique consistent irreducible $\mathbb{Z}$ -gradation $E(5, 10) = \bigoplus_{j \geq -2} p_j$. It is defined by:

$$\deg x_i = 2 = -\deg \partial_i, \quad \deg d_{ij} = -1.$$

One has: $p_0 \simeq \mathfrak{sl}_5(\mathbb{C})$ and the p_0 -modules occurring in the L_- part are:

$$\begin{aligned} p_{-1} &= \langle d_{ij} \mid i, j = 1, \dots, 5 \rangle \simeq \Lambda^2 \mathbb{C}^5, \\ p_{-2} &= \langle \partial_i \mid i = 1, \dots, 5 \rangle \simeq \mathbb{C}^{5*}. \end{aligned}$$

Recall also that p_1 consist of closed 2-forms with linear coefficients, that p_1 is an irreducible p_0 -module and $p_j = p_1^j$ for $j \geq 1$.

We take for the Borel subalgebra of p_0 the subalgebra of the vector fields $\langle \sum_{i \leq j} a_{ij} x_i \partial_j \mid a_{ij} \in \mathbb{C}, \text{tr}(a_{ij}) = 0 \rangle$, and denote by $F(m_1, m_2, m_3, m_4)$ the finite-dimensional irreducible p_0 -module with highest weight (m_1, m_2, m_3, m_4) . We let

$$M(m_1, m_2, m_3, m_4) = M(F(m_1, m_2, m_3, m_4))$$

denote the corresponding generalized Verma module over $E(5, 10)$.

Conjecture 4.1. *The following is a complete list of generalized Verma modules over $E(5, 10)$ ($m, n \in \mathbb{Z}_+$):*

$$M(m, n, 0, 0), \quad M(0, 0, m, n) \quad \text{and} \quad M(m, 0, 0, n).$$

In this section we construct three complexes of generalized $E(5, 10)$ Verma modules which shows, in particular, that all modules from the list given by Conjecture 4.1 are degenerate. We let:

$$S_A = S(\mathbb{C}^5 + \Lambda^2 \mathbb{C}^5), \quad S_B = S(\mathbb{C}^{5*} + \Lambda^2 \mathbb{C}^{5*}), \quad S_C = S(\mathbb{C}^5 + \mathbb{C}^{5*}).$$

Denote by x_i ($i = 1, \dots, 5$) the standard basis of $\mathbb{C}^5$ and by $x_{ij} = -x_{ji}$ ($i, j = 1, \dots, 5$) the standard basis of $\Lambda^2 \mathbb{C}^5$. Let x_i^* and $x_{ij}^* = -x_{ji}^*$ be the dual bases of $\mathbb{C}^{5*}$ and $\Lambda^2 \mathbb{C}^{5*}$, respectively. Then S_A is the polynomial algebra in 15 indeterminates x_i and x_{ij} , S_B is the

polynomial algebra in 15 indeterminates x_i^* and x_{ij}^* and S_C is the polynomial algebra in 10 indeterminates x_i and x_i^* .

Given two irreducible p_0 -modules E and F , we denote by $(E \otimes F)_{\text{high}}$ the highest irreducible component of the p_0 -module $E \otimes F$. If $E = \oplus_i E_i$ and $F = \oplus_j F_j$ are direct sums of irreducible p_0 -modules, we let $(E \otimes F)_{\text{high}} = \oplus_{i,j} (E_i \otimes F_j)_{\text{high}}$. If E and F are again irreducible p_0 -modules, then $S(E \oplus F) = \oplus_{m,n \in \mathbb{Z}_+} S^m E \otimes S^n F$, and we let $S_{\text{high}}(E \oplus F) = \oplus_{m,n \in \mathbb{Z}_+} (S^m E \otimes S^n F)_{\text{high}}$. We also denote by $S_{\text{low}}(E \oplus F)$ the complement to $S_{\text{high}}(E \oplus F)$.

It is easy to see that we have as p_0 -modules:

$$\begin{aligned} S_{A,\text{high}} &\simeq \oplus_{m,n \in \mathbb{Z}_+} F(m, n, 0, 0), \\ S_{B,\text{high}} &\simeq \oplus_{m,n \in \mathbb{Z}_+} F(0, 0, m, n), \\ S_{C,\text{high}} &\simeq \oplus_{m,n \in \mathbb{Z}_+} F(m, 0, 0, n). \end{aligned}$$

Introduce the following operators on the spaces $M(\text{Hom}(S_X, S_X))$, $X = A, B$ or C :

$$\nabla_X = \sum_{i,j=1}^5 d_{ij} \otimes \theta_{ij}^X,$$

where

$$\theta_{ij}^A = \frac{d}{dx_{ij}}, \theta_{ij}^B = x_{ij}^*, \theta_{ij}^C = x_i^* \frac{d}{dx_j} - x_j^* \frac{d}{dx_i}.$$

It is immediate to see that $p_0 \cdot \nabla_X = 0$. In order to apply Proposition 1.1, we need to check that

$$(4.1) \quad p_1 \cdot \nabla_X = 0.$$

This is indeed true in the case $X = C$, but it is not true in the cases $X = A$ and B . In fact (4.1) applied to $f \in S_X$, $X = A$ or B , is equivalent to the following equations, respectively ($a, b, c, d = 1, \dots, 5$):

$$(4.2) \quad \left(\frac{d}{dx_{ab}} \frac{d}{dx_{cd}} - \frac{d}{dx_{ac}} \frac{d}{dx_{bd}} + \frac{d}{dx_{ad}} \frac{d}{dx_{bc}} \right) f = 0,$$

$$(4.3) \quad (x_{ab}^* x_{cd}^* - x_{ac}^* x_{bd}^* + x_{ad}^* x_{bc}^*) f = 0.$$

It is not difficult to check the following lemma.

Lemma 4.2. (a) The subspace of S_A defined by equations (4.2) is $S_{A,\text{high}}$.

(b) Equations (4.3) hold in $S_B/S_{B,\text{low}}$.

(c) Equation $\nabla_X^2 = 0$ is equivalent to the system of equations ($a, b, c, d = 1, \dots, 5$):

$$\theta_{ab}\theta_{cd} - \theta_{ac}\theta_{bd} + \theta_{ad}\theta_{bc} = 0.$$

We let:

$$V_A = S_{A,\text{high}}, V_B(\text{resp. } C) = S_B(\text{resp. } C)/S_{B(\text{resp. } C),\text{low}}$$

The above discussion implies

Proposition 4.3. (a) The operators ∇_X define $E(5, 10)$ -morphisms $M(V_X) \rightarrow M(V_X)$ ($X = A, B$ or C).

(b) $\nabla_X^2 = 0$ ($X = A, B$ or C).

(c) $\nabla_X = 0$ iff $X = A$ and $n = 0$, or $X = C$ and $m = 0$.

The non-zero maps ∇_X are illustrated in Figure 2. The nodes in the quadrants A , B and C represent generalized Verma modules $M(m, n, 0, 0)$, $M(0, 0, m, n)$ and $M(m, 0, 0, n)$, respectively. The arrows represent the $E(5, 10)$ -morphisms ∇_X , $X = A, B$ or C in the respective quadrants.




FIGURE 2.

The following theorem summarizes our results on degenerate $E(5, 10)$ -modules.

Theorem 4.4. (a) *Each connected component in Figure 2 is a complex of $E(5, 10)$ -modules.*
 (b) *The kernels of all morphisms in these complexes are irreducible maximal submodules.*

References

- [CK] Cheng, S-J., Kac, V.: Structure of some $\mathbb{Z}$ -graded Lie superalgebras of vector fields. *Transformation Groups*, **4** (1999), 219-272.
- [K] Kac, V.: Classification of infinite-dimensional simple linearly compact Lie superalgebras. *Advances in Math.* **139** (1998), 1-55.

- [KR1] Kac, V., Rudakov, A.: Representations of the exceptional Lie superalgebra $E(3, 6)$ I: Degeneracy conditions, ESI preprint no 921, 2000.
- [KR2] Kac, V., Rudakov, A.: Representations of the exceptional Lie superalgebra $E(3, 6)$ III: Four series of degenerate modules, *Comm. Math. Phys.* 222 (2001), 611-661.
- [KR3] Kac, V., Rudakov, A.: Representations of the exceptional Lie superalgebra $E(3, 6)$ III: Classification of singular vectors (in preparation).
- [M] Mac Lane, S.: Homology (3-rd corr.printing). Springer-Verlag: NY-Heidelberg-Berlin, 1975.
- [R] Rudakov, A.: Irreducible representations of infinite-dimensional Lie algebras of Cartan type. *Math.USSR-Izvestia* 8 (1974), 836-866.

Authors' addresses:

Department of Mathematics, MIT, Cambridge MA 02139, USA
email: kac@math.mit.edu

Department of Mathematics, NTNU, Gløshaugen, N-7491 Trondheim, Norway
email: rudakov@math.ntnu.no